

The weather today rocks or sucks for my tree: Exploring the understanding of climate impacts on forests at high school level through tweets

Thomas Mölg^{1*}, Jan. C. Schubert^{2*}, Annette Debel¹, Steffen Höhnle², Kathy Steppe³, Sibille Wehrmann¹, Achim Bräuning¹

** These authors contributed equally to this work*

¹Institute of Geography, Friedrich-Alexander-University Erlangen-Nürnberg (FAU), 91058 Erlangen, Germany

²Lehrstuhl für Didaktik der Geographie, Friedrich-Alexander-University Erlangen-Nürnberg (FAU), 90478 Nürnberg, Germany

³Laboratory of Plant Ecology, Faculty of Bioscience Engineering, Ghent University, 9000 Ghent, Belgium

Correspondence to: thomas.moelg@fau.de or jan.christoph.schubert@fau.de

Abstract. With the progression of global warming, impacts on the human sphere will undoubtedly increase. One prominent example at mid latitudes is the stress of forests under climate change, which the project "BayTreeNet" (<https://baytreenet.de/>) addresses from an interdisciplinary viewpoint. Scientists from physical climatology, dendroecology, and educational research collaborate to examine how long-term changes in weather patterns affect the state of trees, and how the atmosphere/tree relation can be used to the advantage of improving the communication of climate change effects to, in particular, high school students. This article presents a 1-week case study for summer 2021, when a distinct variability in weather patterns induced significant tree responses. The students of seven selected partner schools commented on the measured tree and weather data, which were available in real-time, in the form of tweets and in conjunction with their educational geography program. The analyses of the tweets reveal the result that the students succeed in verbalizing the measured weather and, furthermore, manage to draw linkages to the stem diameter changes of trees. Problems arise with the use of less perceivable variables like the sap flow in the trees; also, the student posts exhibit shortcomings in establishing causal connections. Hence, the case study points to a discrepancy between describing basic environmental information and appreciating, or understanding, the underlying mechanistic links. This point will serve to refine future classroom concepts and, moreover, to enhance the communication of climate change effects on forests for the general public.

1 Introduction and general motivation

With the arrival of evening, TV prime time approaches and one show with it – the daily weather forecast. “The world’s most watched science show” is, therefore, a statement that has appeared in the public and in the media. The implied importance of weather news has even increased with the rise of new, internet-based technologies over the past 20 years, which have led to a multiplication of weather information services. A related scientific study for the U.S. concluded, unsurprisingly, that "weather forecasts are a daily part of the lives of the vast majority of the U.S. public" (Lazo et al., 2009, p. 775). The roots of this circumstance go far back in time – let’s think of Aristoteles’ treatise "Meteorologica" or the century-long association of weather with the divine, which reflect the general desire of humans to understand weather. The side effect of this deeply-anchored interest is a familiarity of the public with

37 some central (scientific) terms like *high pressure*, *front*, and *cyclone*, and a sense of what they mean in practice
38 for local weather phenomena, or generally speaking for impacts (e.g., on one's own daily life, on health, or on the
39 environment). The scientific synthesis of these terms is expressed in the systematic study of daily large-scale
40 weather patterns in the atmosphere, which was mostly developed for central Europe in the 1940s/50s, and has to
41 become known by its German term "Großwetterlage" (see, for example, James (2007) for a short historical
42 overview). Although the scientific concept as well as the daily consumption of the topic in public media refer to
43 *weather*, meaning the state of the atmosphere ranging from a short moment in time to a few days, there is also a
44 *climate* dimension. Consider that certain weather patterns become less or more frequent over decades and longer.
45 The consequences will be a change in the seasonal or annual characteristics of air flow patterns (which we call the
46 large-scale *atmospheric circulation*), and hence a change of the surface climate at locations in the reach of the
47 weather patterns. This relation is the starting point for connecting weather patterns to our changing climate, which
48 has been studied extensively in recent research (see Sect. 2). Importantly, the scientific weather pattern term mostly
49 encapsulates the distribution of air pressure centres and the resultant large-scale circulation modes. These
50 atmospheric dynamics are also embodied by the elements that appear on weather maps in the daily news or on our
51 cell phones as "H" or "L" and wind arrows, feeding the public with the basics of an important climate change
52 aspect. The present article originates from a transdisciplinary project that aims to exploit the omnipresence of
53 weather information in today's world, and the increasing public awareness of climate change impacts on the
54 environment, ecosystems, and human health.

55 In this context, a prominent impact case is the response of forests to climate change. Trees are important organisms
56 for forest ecosystems and society as they provide ecological services, and tree-ring research (dendrochronology)
57 is a well-established discipline to assess the impacts of climate on tree growth and to reconstruct climate conditions
58 of the past. Although a number of concepts and tools have been developed for introducing the basics of tree-ring
59 research in different formats to the public, such as articles or videos (e.g., Davi et al., 2022), there is hardly any
60 tool to monitor the effects of weather on tree performance in real time, and to get the relevant explanations of the
61 climatic background. Here, we target this gap by making real-time weather and tree data available to high schools
62 in conjunction with their geography programs, in order to investigate characteristics of climate change
63 communication and education at the high school level. While this provides the general motivation, Sect. 2 will
64 explain the specific circumstances as well as the structure of the research and the article.

65

66 **2 Specific context and goals**

67 The transdisciplinary project mentioned is called *BayTreeNet* and was described in a recent publication from a
68 conceptual viewpoint (Bräuning et al., 2022). The project was established in response to the evidence of increasing
69 drought stress on forests in Bavaria in the most recent decades (Debel et al., 2021) and, hence, presents an effort
70 to generate more knowledge of this threat with regard to future climates. *BayTreeNet* tackles the topic from the
71 perspectives of three special fields. Physical climatology, dendroecology, and educational research collaborate
72 under the overarching goal of increasing public awareness and preparedness for stress on forest ecosystems under
73 climate change, which must build on a solid communication and education of the problem. Besides the
74 transdisciplinary basis, a novel facet of *BayTreeNet* is to look at the climate/forest interaction at the scale of

75 weather patterns (henceforth GWL, according to the German term "Großwetterlage" introduced in Sect. 1), which
76 adds value to the traditional approach of relating forest responses to more aggregated climate data (e.g., monthly
77 or annual mean values). The GWL resolution, in turn, provides the connecting link to the public weather
78 information (Sect. 1), and is therefore a promising starting point for enhancing dissemination and education in the
79 given context.

80

81 In the physical climatology sub-project, GWL in the past and in the future and their manifestation at the local level
82 are examined with the help of meteorological observations and climate modelling. An important initial result was
83 the generation of a new climate data set for Bavaria (1987-2018) with a very high resolution in space (Collier and
84 Mölg, 2020). The dendroecology sub-project relies on a network of measurement sites over Bavaria, where
85 important tree growth variables are monitored (Sect. 3.1). A first study using these data concluded that the tree
86 growth/climate relationships show an elevation dependency and, moreover, have not been constant over the past
87 five decades (Debel et al., 2021). The core of the educational sub-project, in turn, is a collaboration with high
88 schools in the vicinity of the measurement sites. Their students (grades 10 and 11 and 15-17 years old) observed
89 the weather and tree growth data in real-time, which is enabled by an accessible interface on the project webpage
90 (<https://baytreenet.de/>), and subsequently translated their interpretation of the trees' physiological responses to
91 simple Twitter (now known as X) messages. This translation embodies the educational aspect and provides the
92 data for the communication research of the present study.

93

94 Individually, the three mentioned scientific fields can build on a rich literature with regard to the present topic.
95 Investigations in climatology have, for instance, targeted the role of GWL for health, extreme events, hydrological
96 and ecological processes (Bissolli, 2001; Post, 2002; Clark and Brown, 2013; Loikith et al., 2017; Pisistaki et al.,
97 2020; Zong et al. 2022). Given the concept's history (Sect. 1), central Europe has always been a focus (e.g.,
98 Riediger and Gratzki, 2014; Herrera-Lormendez et al., 2021). In dendroecology, studies of the dependence of tree
99 growth on climatic variables in central Europe have an equally long history; some major recent focus points were,
100 for example, how summer temperature and rainfall variability affect the growth of European tree species (Debel
101 et al. 2021; Dulamsuren et al. 2017; Friedrichs et al. 2009; Kraus et al. 2016). And finally, contributions from the
102 field of educational research on climate change deal, among other things, with the incorrect or insufficient ideas
103 of students. For example, several conceptual models of the causes of the anthropogenic greenhouse effect have
104 been identified among students, which contradict scientific explanations (including the wrong explanation of the
105 greenhouse effect via the hole in the ozone layer) and can hinder learning processes (Choi et al. 2010). With respect
106 to consequences of climate change, it was shown that students can name a variety of examples but have major
107 problems with spatial differentiation (Boyes, Stanisstreet 1993; Leiserowitz 2021). Similar results were shown to
108 concern many adults as well (Dove 1996).

109

110 However, to our knowledge no scientific project has brought together the three disciplines for the purpose of
111 fostering communication and education, regarding the important topic of forest response to climate change. In the
112 present article, we make a first effort to synthesize the multidisciplinary data through a 1-week case study for early
113 summer 2021. The main goal is to examine what level of understanding exists in high school students about the
114 forest/atmosphere interactions, by considering the associated Twitter data. Our working hypothesis is that students
115 are basically successful at translating weather data into words (henceforth called "verbalizing"), yet more

116 difficulties exist with verbalizing tree data and, in particular, with correctly describing the relationships between
 117 weather and tree data. This hypothesis is derived from the considerations that weather reports and conversations
 118 about weather are a part of the students' everyday life (cf. Sect. 1), while the terms related to the state of trees are
 119 much less known, so students have less practice in verbalizing information on tree growth. Hence, linking weather
 120 and tree data represents the greatest challenge and involves a comparatively high complexity.

121
 122 To realize our approach, we selected a week with obvious variability in weather patterns and tree responses, and
 123 at the same time, with a relatively high number of student tweets. Since identifying such a week involved screening
 124 the physical-science data as well, we first present the climatological and dendroecological input from our project
 125 in Sect. 3; this is a preparatory step that is needed to comprehend the entire study. The actual communication
 126 (educational) research is then presented in Sect. 4, before the conclusions in Sect. 5.

127 **3 Physical science basis**

128 **3.1 Data and methods**

129 **Weather pattern classification.** In climatology, daily values of sea-level pressure and geopotential height at 500
 130 hPa (roughly, the mid-troposphere level) are the typical input variables for the classification of GWL. The most
 131 common and popular classification was introduced and published by Hess and Brezowsky in 1952 in the Catalogue
 132 of GWL in Europe (see James, 2007), containing daily weather patterns since 1881. The national German
 133 Meteorological Service (DWD) has been using this classification since then. The manual classification of Hess
 134 and Brezowsky yields 29 types of GWL, which are characterized by the air pressure pattern, cyclonicity, the main
 135 direction of large-scale air flows, and vorticity (Werner and Gerstengarbe 2010). The distribution, intensity, and
 136 characteristics of high- and low-pressure systems define the different types of GWL, which need to last at least
 137 three consecutive days. GWL types show different occurrence probabilities during the year and vary among the
 138 seasons, and each GWL can bring significantly different weather to certain regions and locations in Europe. Table
 139 1 summarizes some relevant aspects of the traditional GWL scheme, which is the basis for the present study.

140
 141 **Table 1:** Some examples of the 29 GWL (frequent ones and relevant ones for our study) in original German and
 142 translated into English definitions after James (2007) and their mean annual probabilities (1881-2008) after Werner
 143 and Gerstengarbe (2010).

144

GWL	Original definition (German)	Translated definition (English)	Probability
WZ	Westlage, zyklonal	Cyclonic Westerly	15.70 %
HM	Hoch Mitteleuropa	High over Central Europe	8.89 %
BM	Hochdruckbrücke (Rücken) Mitteleuropa	Zonal Ridge across Central Europe	7.72 %
NEa	Nordostlage, antizyklonal	Anticyclonic North-Easterly	2.18 %
HNFa	Hoch Nordmeer-Fennoskandien, antizyklonal	High Scandinavia-Iceland, Ridge C. Europe	1.14 %

145
 146 **Wood formation detection.** In the dendroecology sub-project, we have established a dendroecological network
 147 of eleven study sites (initially ten) to monitor responses of forest ecosystems to changing climate dynamics at

148 different temporal resolutions. While long-term growth changes, adaptations of the wood anatomical structure and
149 adaptation of intrinsic water use efficiency are assessed at decadal to annual time scales, cambial growth dynamics
150 and stem diameter variations are monitored at interannual to even hourly resolution. While these dendroecological
151 sample sites are located inside forests, an additional monitoring method detects tree responses to changes in the
152 environment in real-time. For this purpose, single tree individuals close to each sample site were equipped with an
153 internet-based logger and sensors to measure sap flow and stem diameter variations. The sap flow sensor measures
154 the sap flow transport rates ($\text{cm}^3 \text{hour}^{-1}$) from the roots to the crown (Smith and Allen, 1996; Steppe et al., 2010,
155 2015; Vandegehuchte and Steppe, 2013). The point dendrometer registers stem diameter changes (mm) caused by
156 the reversible shrinking and swelling of living stem cells in 20 minute intervals. These are visible as daily cyclic
157 stem diameter variations, the amplitudes of which inform us about the water status of the tree (Deslauriers et al.
158 2007; Drew and Downes 2009; Steppe et al., 2015). In addition, the dendrometer records the irreversible growth
159 of the xylem and phloem by the formation of new xylem (wood) and phloem (bark) cells, which becomes visible
160 as a long-term trend of increasing stem diameter superimposed on the daily radial variations. While under humid
161 conditions, the long-term trend is positive, tree stems may shrink in diameter to a certain extent under drought
162 conditions. The mentioned sensors were developed by the research group at Ghent for the initiative
163 “TreeWatch.net”, details of which can be found in Steppe et al. (2016).

164


165 **Data preparation.** These two high-precision instruments indicate the individual trees’ responses to varying GWL
166 and the associated weather conditions like, for example, a heavy rain event or a heatwave. The measured data are
167 transmitted real-time to the PhytoSense Cloud service, which is a tool for data storage, analysis, processing, and
168 running model simulations to visualize the trees’ hydraulic function and carbon status (Steppe et al., 2016). Finally,
169 the processed data are sent to the project homepage (<https://baytreenet.de/>), where the physiological data of each
170 “talking tree” are graphically displayed. At the same time, information about the current weather at each site, e.g.,
171 precipitation and temperature, is available on the homepage. This includes a map of central Europe with the current
172 flow patterns at the 500 hPa (~5 km) and 850 hPa (~1.5 km) geopotential height levels, and a weather map
173 indicating the location of fronts. Therefore, the short-term responses of trees to weather and long-time tree growth
174 become detectable and displayable.

175 **3.2 Results: Weather variability and tree response**

176 The calendar week 22 in early summer 2021 (May 31st to June 6th) was detected appropriate for this case study.
177 In this week, two GWL were present: NEa (Anticyclonic North-Easterly) on Monday, Saturday, and Sunday (May
178 31st, June 5th and June 6th) and HNFa (High Scandinavia-Iceland, Ridge Central Europe) from Tuesday to Friday
179 (June 1st to June 4th). Both GWL are typical weather types for the early summer season and show their maximum
180 likelihood in May (HNFa) and June (NEa). Measured against the full year they occur with a probability of 1.41 %
181 (NEa) and 2.18 % (NEa) (cf. Table 1). The two GWL belong to anticyclonic types and typically induced the highest
182 average air temperatures in the recent past. For example, HNFa shows a deviation of +4.36 °C in the daily mean
183 air temperature with reference to the period 1951-1978 (Werner and Gerstengarbe 2010). Considering
184 precipitation, both GWL coincide with drier conditions on average, however, the drying signal is not as significant
185 as the warming signal.

186

187 These average anomalies are consistent with the air temperature characteristics in June 2021, as seen in the
 188 measurements at the official DWD weather stations. In this month, particularly high temperatures were recorded
 189 throughout Germany and daily anomalies of up to +4.5°C (compared to the 1961-1990 mean) occurred locally.
 190 For Southern Germany the situation was very similar. In the case study week, the air temperature anomaly
 191 increased over time until June 3rd and June 4th, when it reached the maximum of +4.9°C departure (same reference
 192 period), and subsequently decreased until the end of the week. These characteristics are also reflected in four
 193 stations close to our tree sites (Fig. 1), which were selected on the basis of reliable tree data and sufficient Twitter
 194 messages. Precipitation at the 35 DWD weather stations in Southern Germany shows a steep increase from June
 195 3rd with its maximum on June 5th and a subsequent decrease on June 6th. At three of the four sites investigated,
 196 a precipitation event is recorded on June 5th with values up to 20 mm (Wunsiedel), while at the fourth site
 197 (Veitshöchheim) this occurs one day later with more than 30 mm (Figure 1).
 198
 199 To summarize the case study week, strong positive air temperature anomalies were measured in Southern Germany
 200 and Bavaria (after cold phases in May), reaching their maximum roughly in the middle of the week. Shortly after
 201 this temperature maximum, precipitation occurred after several days of drought. Thus, the selected 1-week window
 202 was characterized by a transition to warmer and wetter conditions.
 203




204
 205 **Figure 1:** Two-meter air temperature (dot symbols) and precipitation (rectangles) at four selected weather stations
 206 in Bavaria during the calendar week 22 of the year 2021 (data basis: DWD).
 207

208 How did these atmospheric conditions affect the sample trees? In general, most talking trees showed similar
 209 responses to the prevalent weather. The tree responses only differed in the timing between different locations,
 210 since the rain events happened asynchronously between sites, as discussed above.
 211

212 From June 1st to 4th, the influence of the HNFa weather pattern (with its dry and warm weather conditions) led to
 213 characteristic tree physiological response patterns (e.g., Steppe et al., 2015) that indicate sufficient water supply
 214 without water shortage (Fig. 2). Initiated by transpiration and regulated by the stomata, the water transport from
 215 the roots to the crown started shortly after sunrise. Consequently, the sap flow rate increased rapidly in the early
 216 morning, peaked around noon, and remained at high level until the evening. After sunset, the sap flow rate dropped
 217 to pre-dawn level, showing minimum values in the early morning hours before the beginning of the new day's
 218 cycle.

219
 220 The dendrometers recorded daily cycles of diameter changes, mainly due to the reversible shrinking and swelling
 221 of the stem caused by the depletion and replenishment of the stem water stored in wood and bark tissues. Stem
 222 diameters decreased in the morning shortly after sunrise, since photosynthesis started and water transpired through
 223 the open stomata. As long as transpiration persisted and water flowed from the roots to the crown, the water
 224 reserves in the stem depleted and the stem diameter shrank. After sunset, stem diameter began to increase again
 225 due to the cessation of transpiration and the refilling of water reserves in the stem (Steppe et al., 2006). Beside
 226 these daily cycles, the maximum stem diameters increased from day to day because of irreversible stem growth
 227 driven by turgor and formation of new xylem cells.

228
 229 With the occurrence of rather strong rain events later in the week, both data series show a significant change (Fig.
 230 2). On the one hand, the sap flow rate of most (7 out of 10) trees did not increase to the same degree as in the days
 231 before, and furthermore, the sap flow of all talking trees decreased earlier around midday. A high cloud cover
 232 fraction and high air humidity reduced transpiration, hence the stem water transport declined during and after the
 233 rain event. On the other hand, the talking trees overwhelmingly showed a sharp increase in diameter shortly after
 234 the rainfall. Moreover, the diameter curve showed no decline during the daytime, pointing to continuous refilling
 235 of the internal water storage pools that were emptied during the preceding dry period. Thus, higher air humidity
 236 and less vertical water transport caused intensified water storage in stem cells. As a result, the cells got water-
 237 saturated, and the depletion effect (causing a decrease in diameter) was mitigated.



238
 239 **Figure 2:** Stem diameter variations and sap flow rates of *Tilia platyphyllos* at Burghausen as an example of tree
 240 response during the period May 31st to June 6th, 2021. Note that from June 5th to 6th, precipitation occurred,
 241 leading to a strong increase in stem diameter.

242 **4 Communication and educational research**

243 **4.1 Methodology**

244 We developed a half-day introductory workshop for the participating students and held it on site at each school
245 (~200 students at 10 schools initially). When the Covid19 pandemic began, we converted the workshop into a self-
246 study course that could be completed online. The participating groups of students changed every school year; each
247 group took part in the course at the beginning. In terms of content, the course aimed to build up a basic
248 understanding of GWL as well as of tree growth conditions, as these aspects play a marginal role in the curriculum
249 and, therefore, there was little previous knowledge to refer to. In addition, exercises were included on how to
250 interpret tree data, in particular, what does (high/low) sap flow mean and what is meant by the variation in stem
251 diameter of a tree? Together with the available information on current weather patterns, the students at each
252 participating school were asked to translate the information on the tree's state as well as the connections with the
253 weather situation into linguistic messages in the form of tweets: short, concise, in everyday language. To practise
254 this, exercises and examples (using current data) were included in the introductory workshop.

255
256 The participating schools dealt with the challenge of organizing the work with the talking trees beyond the regular
257 (curriculum-related) classes in different ways, depending on the local situation. For this reason, the mentoring of
258 the students after the end of the introductory workshop as well as the organization (e.g. when which students at
259 the respective school took over the tweeting) was carried out by supervising teachers at each school. At some
260 schools, a selected class was responsible for tweeting, while at other schools a group of voluntary students from
261 different classes formed to write the tweets. In addition, at some places individual students were responsible for
262 the tweets for a short period of time and there was a change after two weeks, while at other schools individual
263 students tweeted for much longer. In this context, it should be noted that we asked the students to tweet as regularly
264 as possible. As the school year progressed, the mentoring teachers tried to motivate the students by reminding
265 them to tweet and pointing out the importance. However, it was always the case that tweeting was an additional
266 activity that was not part of the regular lessons, was not included in the grading and was voluntary. Many students
267 wrote the tweets in their free time, and tweeting continued at many schools even during school vacations.

268
269 The fundamental decision to use Twitter in this project was based on several reasons. Firstly, the central task was
270 to convert data into language, which fits well with Twitter as a primarily language-based medium. Secondly, the
271 limited number of characters per message that Twitter had implemented at the time of the project required concise
272 writing with a focus on the central aspects. At the same time, Twitter enables simple, low-threshold communication
273 with each other, which fitted in well with the idea of letting the trees communicate with each other. In addition, it
274 seemed easier to attract the attention of a wider public via Twitter than via conventional websites. Other social
275 media such as TikTok would probably have been closer to the students' everyday lives, but the focus on visual or
276 auditory content would have been less suitable for the project's objectives.

277
278 For the present case study, we analyzed the students' tweets during the carefully selected week in early summer
279 2021 (Sect. 3) with one guiding question: To what extent are the tweets (as linguistic translations of tree data and
280 weather data) appropriate to the subject? The methodological procedure for evaluating the tweets involved several
281 steps. As part of the preparation for the data analysis, the tree data as well as the tweets for all sites were reviewed,

282 and sites where central data were missing were excluded. For the investigated period, one site was excluded due
283 to missing tree data (Bad Reichenhall, sensor error) and another site where no tweets were made (Ettal). Another
284 site (Immenstadt) could not be considered because tweets were only available for three days and, in addition, the
285 tree data provided only limited meaningful information as the tree on site was dying.

286

287 For the remaining seven sites, the weather data and the tree data were prepared in the form of diagrams (e.g., Sect.
288 3) and the students tweets were converted into an editable version suitable for coding. Emojis that did not influence
289 the statement but were merely illustrative in nature were not considered in this process. Several evaluation or
290 analysis steps followed, oriented along the principles of a “thematic analysis” (Braun and Clarke, 2006):

291 (1) Becoming familiar with the data: Through repeated and close reading of the tweets, also against the background
292 of weather data and tree data (as described in Sect. 3.2), an overview and a deeper understanding of the tweets
293 were created by the researchers. This also involved description of the frequencies of tweets per site and per day.

294 (2) Coding with initial codes: The data material was coded according to five deductive categories. These included
295 statements about (i) weather, (ii) growth/stem diameter, and (iii) sap flow. In addition, statements in which students
296 (iv) combined two or more of these categories (for example, effects of weather on tree growth/stem diameter) were
297 coded in a separate category. The same was true for (v) statements in which there was an exchange between
298 students from different tree sites.

299 (3) Search for patterns and salient features in the categorized data: Within the categories, statements were screened
300 for characteristic commonalities or patterns. In addition, conspicuous features were examined. The guiding
301 principle here was a comparison with the scientific basis, i.e. the weather data and tree data in the form available
302 to the students on our homepage. This interpretative step was initially undertaken independently by two
303 researchers, who then compared the interpretations and attempted to reach a consensual interpretation in the event
304 of discrepancies. The results of this step were, in turn, presented and discussed in the working group, which is
305 linked to the goal of reducing subjectivity that is present in interpretative steps.

306 (4) The results of the analyses and interpretation were summarized, written down and supplemented in many places
307 with illustrative quotes from the tweets (see Sect. 4.2).

308 **4.2 Results: Interpretations by high school students**

309 The initial analysis of the tweets showed that three of the seven schools had at least one tweet for each of the seven
310 days of the study period. For the four other schools, tweets were not available for every day, but for at least five
311 days of the study period. In total, the data corpus included 43 tweets (Tab. 2). All 43 of the tweets contained
312 statements about growth or stem diameter changes, 39 contained statements about sap flow, and 37 contained
313 statements about weather. In 13 tweets, connections were made between at least two of the three elements. A
314 sample tweet is shown in Fig. 3.

315

316 **Table 2:** Overview of tree locations and tweets.

317

Locations of the "Talking Tree"	Number of tweets	Number of days with tweets
(1) Neuschönau	7	7
(2) Bad Neustadt	7	6

(3) Veitshöchheim	7	7
(4) Wunsiedel	7	7
(5) Tennenlohe	5	5
(6) Burglengenfeld	5	5
(7) Burghausen	5	5

318



319

320

321 **Figure 3:** Sample tweet for illustrative purposes at the location Neuschönau, meaning “Hello. As hoped for
 322 yesterday, today was great weather, sunshine all day. My diameter has grown by quite a bit and my sap flow has
 323 also increased. I hope you had a great day like me.” (BayTreeNet Neuschönau, 2021).

324

325 The following conspicuous features and characteristics were identified during the more in-depth analyses of the
 326 tweets.

327

328 **(1) Statements about the weather.** The statements about the weather are almost universally in line with the
 329 weather data; even the precipitation events that occurred after previous sunshine are correctly recognized and
 330 named (“Today was another sunny day” [Heute war wieder ein sonniger Tag] // “In any case, it rained and
 331 thundered for all it was worth at my place” [Bei mir hat es jedenfalls geregnet und gewittert, was das Zeug hält],
 332 Wunsiedel). In general, with regard to the weather, it can be seen that not all weather elements that are recognizable
 333 for the students are consistently addressed in the tweets and, moreover, relevant for the tree responses. Regarding
 334 basic weather elements, references to sunshine and cloud cover, to dryness and precipitation, and to cold and
 335 warmth would have been possible. In the tweets, however, predominantly only one of these elements was
 336 addressed. Only in individual cases at least two weather elements were cited together (“Since it was pleasantly
 337 warm again today, I was in a good mood despite Monday, even though it rained today and it was quite cloudy.”,
 338 Tennenlohe). In addition, at some locations, different weather elements were mentioned when comparing the days.
 339 Thus, on some days there were comments (only) about the temperature, on other days (only) about the sunshine.

340

341 **(2) Statements on stem diameter.** The growth of the trees or changes in the stem diameter were consistently taken
 342 up in all tweets. The central common feature of most tweets is that a statement was made about the (thickness)
 343 growth of the tree stem or, more generally, that the tree had grown. However, the recognition or interpretation of
 344 the diurnal fluctuations of the stem diameter seems to be problematic. With very few exceptions (“Well, my
 345 diameter always goes steeply downhill during the course of the day, but I’ve grown a bit anyway” [Also so im
 346 Laufe des Tages geht es ja mit meinem Durchmesser immer steil bergab, aber ich bin trotzdem ein bisschen
 347 gewachsen], Veitshöchheim), these fluctuations were not discussed at all. Instead, a single value of the stem

348 diameter was often compared with a single value from the previous day. In some cases, there were incorrect
349 interpretations from a scientific point of view, because it was not the minimum and maximum stem diameter of
350 one day that were compared with those of the previous or next day, but individual values at different times of the
351 day.

352

353 **(3) Statements on sap flow.** There are a similar number of tweets on sap flow as on weather. It is noticeable that
354 the sap flow was more often described in a relatively vague and superficial way compared to the weather and the
355 stem diameter. A typical example of what was frequently found is "The sap flow is good" [Der Sapflow ist gut].
356 With very few exceptions ("My sap flow was greater today at noon than yesterday, only dropping over the
357 afternoon" [Mein Saftfluss war heute Mittag größer als gestern, nur über den Nachmittag hin ist er gefallen];
358 Neuschönau), diurnal variations in sap flow were not picked up in the tweets. Instead, individual values or
359 maximum values of different days were compared (similar to stem diameter).

360

361 **(4) Links between weather, stem diameter and sap flow.** Linkages in the sense of causal relationships between
362 weather, growth/stem diameter, and sap flow were rarely the subject of tweets. In most cases, the information on
363 the three areas was rather unrelated to each other ("Hey friends. Today was finally awesome weather. My sap flow
364 is shooting through the roof and I'm thinning out a bit. I hope you're all well" [Hey Freunde. Heute war ja mal
365 endlich Bombenwetter. Mein Saftfluss schießt durch die Decke und ich werde ein bisschen dünner. Ich hoffe euch
366 geht's gut], Burglengenfeld // "Hello. As hoped yesterday, today was great weather, sunshine all day. My diameter
367 has grown by quite a bit and my sap flow has also increased" [Hallo. Wie gestern erhofft war heute ein super
368 Wetter, den ganzen Tag Sonnenschein. Mein Durchmesser ist um einiges gewachsen und auch mein Saftfluss ist
369 mehr geworden], Neuschönau). If links were made, the typical pattern was to start with the weather and go to the
370 diameter or the sap flow ("But at least I was able to grow due to the rain" [Aber durch den Regen konnte ich
371 wenigstens wachsen], Burglengenfeld // "My diameter has grown a gigantic amount due to the great weather"
372 [Mein Durchmesser ist durch das tolle Wetter nochmal ein gigantisches Stück gewachsen], Tennenlohe). Links
373 between sap flow and diameter were not made.

374

375 **(5) Communication between the tree sites.** In many cases, the other sites or the other trees were addressed in the
376 salutation or the closing words ("Hello friends" [Servus Freunde], "Hello my dear fellow trees ["Hallöchen meine
377 lieben Mitbäume]" as well as "Well, we'll hear each other tomorrow" [Naja wir hören uns dann morgen],
378 Burghausen). In some cases, questions were also addressed to the other tree sites ("How was everything going
379 with you?" [Wie war es denn bei euch?], Wunsiedel). In contrast, an explicit take-up of such questions as well as
380 of content-related information on tree data and weather data shared by other tree sites occurred with very few
381 exceptions ("Hi Karl. I think your wish for rain came true." [Hi Karl. Ich glaube, dein Wunsch nach Regen ist in
382 Erfüllung gegangen], Wunsiedel). An exchange on measured values on the level of tree data was not detectable at
383 any site.

384

385 In terms of summary and discussion, it can be stated that the students implemented the content of the online
386 introductory course appropriately and well, at least on the formal or surface level. The number of tweets was
387 relatively high (= daily) for most schools during the analysed period, and the tweets commented relatively
388 consistently on weather, stem diameter/tree growth, and sap flow (as the three key metrics). The change in weather

389 on June 5th was recognized and commented on. Thus, the central goal of having students verbalize measured data
390 was achieved. With a detailed look at the tweets as well as in comparison with the tree data and weather data
391 available for the respective location and period, it became clear that the tweets were not comprehensive (weather)
392 or only considered parts of the measurement data (growth, sap flow). In particular, the fluctuations in stem diameter
393 and sap flow over the course of the day, considered individually, seem to be too complex for the students to be
394 addressed in their tweets. The connections between sap flow and stem diameter, which are even more complex,
395 were not addressed by the students at any point. In contrast, weather elements were used (although only in a few
396 instances) to show effects on the trees. This approach is scientifically correct and in line with the students'
397 perceptions: Weather and weather elements are known to students from their own experience and from weather
398 reports, see Sect. 1, and they can also feel them with their own senses. In contrast, changes in stem diameter are
399 not perceptible with their own senses, at least within the resolution that occurred in the case study period. The
400 same applies to the sap flow of trees. Both are only accessible via the sensors and graphical illustrations of the
401 measured values. Starting from the weather, which is familiar and perceptible from everyday life, the students
402 have tried to explain the tree data.

403

404 The goal of initiating a communicative exchange between students from different tree locations was only partially
405 successful. Although there was almost always an address to other locations, the exchange was almost without
406 exception on a superficial level. However, there was almost no exchange or discussion of weather data or
407 measurement data between the sites. One reason could be that even understanding one's own tree data is
408 challenging and not always successful. Thinking through data from other trees would mean an additional increase
409 in demand and effort.

410 **4.3 Limitations**

411 The results presented are subject to a number of limitations. Firstly, this is a case study in which we looked at a
412 limited time period, which is not necessarily characteristic of other times. The decision for the period examined
413 was deliberately made on the basis of weather and tree data, as the situations at the schools were not homogenous
414 (see Sect. 4.1), the tweeting students changed over time, and the tweets could not be assigned to individual students.
415 For this reason, the data basis does not allow for an analysis of learning gains over the course of a school year or
416 for similar analyses, which is why we opted for a case study based on the physical-science observations. So while
417 the tweets can be seen as a random selection to some extent, it remains unclear whether the study period is
418 characteristic of other time periods. In addition, there were technical problems (sensor failure at one site) and not
419 all schools sent out tweets during the week studied.

420

421 Besides the fundamental questions about the significance of Twitter in comparison to other social media in
422 students' everyday lives, the medium Twitter and the associated restriction to a maximum of 140 characters per
423 tweet played a special role. On the one hand, this allowed the measured values to be verbalized by the students in
424 a condensed manner and an accessible one for the interested public. At the same time, discussing profound causal
425 links between tree and weather data as well as a communicative exchange on the measurements between the sites
426 are made more difficult by this format alone. In retrospect, it would have been useful to explicitly encourage
427 students to submit multiple tweets within one day. At the same time, from a realistic perspective, it must be stated

428 that looking at the tree data and tweeting already meant an additional task for the students. This made it difficult
429 to increase the requirements further.

430

431 Overall, therefore, the fact that links between weather, stem diameter, and sap flow, as well as between tree
432 locations, occur very infrequently cannot be clearly or solely attributed to comprehension problems or the like;
433 one reason may also be the format (140 characters). Although in some places it can be assumed that there were
434 indeed difficulties in comprehension, it can be seen as positive that the students succeeded in transferring the tree
435 data into short tweets in everyday language.

436 **5 Conclusions and Outlook**

437 If we close the circle to the introduction of this article (Sect. 1), the 1-week period in summer 2021 was a good
438 choice in the sense that it showed ample weather variability. Warming in the first half was followed by heavy rain
439 events at our study sites, which induced significant tree responses. These were characterized by a growth in stem
440 diameter and a high sap flow rate, while pronounced diurnal cycles were superimposed on these weekly tendencies.
441 In connection with the tweet data, a first idea of how high school students understand the relationship between
442 atmospheric variability and tree responses can be shaped.

443

444 The results demonstrate that students put a focus on weather elements, which underlines the connection of people
445 to daily weather information emphasized in the introduction. Regarding the tree variables, it seems that they pick
446 up the stem diameter changes more easily than sap flow variability. This is not uncommon, and may have several
447 reasons: compared to sap flow, stem diameter is easier to perceive, whereas sap flow is more elaborate and abstract.
448 In addition, trees with trunks of different thicknesses are part of our lifeworld context, presumably more prior
449 knowledge is available for stem changes than for sap flow. New information can be embedded in existing mental
450 models more easily as long as the new information is consistent with the existing mental model. Such an
451 importance of contexts and prior knowledge is well known in cognitive educational science (for example
452 Podschuweit and Bernholt, 2020; Witherby and Carpenter, 2022). One common problem irrespective of the tree
453 variable was the appreciation and understanding of the distinct diurnal cycles in stem diameter change and sap
454 flow. In the authors' experience from teaching at university level, understanding the meaning of such daily cycles
455 (as a form of internal climate variability) appears to be hard in general. Another common problem was that causal
456 relationships between the various variables were hardly made, except some basic characteristics of how weather
457 tendencies affect the tree responses. While this represents without doubt a more complex question, the deficiency
458 also suggests that teaching the climate as a physical cause/effect system could probably be enhanced at school
459 level. Altogether, the results confirm our working hypothesis and point to a discrepancy between verbalizing
460 environmental information and drawing mechanistic links.

461

462 In the near future, it is planned to complete the study of systematic GWL changes over many decades, which will
463 allow the students to relate the topic more strongly to the climate change aspect. We will also attempt to maintain,
464 and even to extend, the tree network in the future, either by a follow-up research project or by other means of
465 public education. For future classroom sessions, teaching materials that focus on the relationship between weather
466 conditions and the responses of trees are currently developed and evaluated. These connections are to be worked

467 out both in general and in concrete case studies. For the latter, exemplary periods with tree data and tweets of the
468 students will be included. In addition, further time periods with relevant weather changes and the corresponding
469 tree reactions will be analyzed in order to better assess the robustness of the findings in this case study.

470

471 **Data availability.** Weather data are openly available from the German Weather Service (DWD). The tree data and
472 associated tweets can be accessed on our website at <https://baytreenet.de/>, by selecting a study site and adjusting
473 the date range to 31 May to 6 June 2021.

474 **Author contributions.** AB, TM and JCS designed the study and arranged the project funding. AD, JCS, SH and
475 SW analyzed the data. KS supported the tree sensor data collection. All authors contributed to the writing and
476 editing of the manuscript, which was coordinated by TM.

477 **Competing interests.** The authors declare that they have no conflict of interest.

478 **Ethical statement.** Ethical approval for this research was given by the Friedrich-Alexander-University (FAU)
479 Erlangen-Nürnberg ethics committee.

480 **Acknowledgements.** This research was funded by the Bavarian State Ministry of Science and Arts, as part of the
481 Bavarian Climate Research Network (bayklif). We thank Katrien Schaepdryver, supported by the BOF research
482 project TreeWatch (grant no 01J07919), and Erik Moerman from Ghent University for their technical support. We
483 also thank the teachers and students of the partner schools for their big engagement in this project. And finally,
484 big thanks to the two reviewers and executive editor John K. Hillier for very helpful comments on earlier versions
485 of this manuscript.

486

487 **References**

488 BayTreeNet Neuschönau [BayTreeNet]: "Hello. As hoped for yesterday, today was great weather, sunshine all
489 day. My diameter has grown by quite a bit and my sap flow has also increased. I hope you had a great day like
490 me." Twitter, available at: <https://x.com/6baytreenet?s=43> (last access: 23 November 2023), posted: 19:30, 2
491 June 2021.

492 Bissolli, P. and Dittmann, E.: The objective weather type classification of the German Weather Service and its
493 possibilities of application to environmental and meteorological investigations, *Meteorol. Z.*, 10, 253–260,
494 doi:10.1127/0941-2948/2001/0010-0253, 2001.

495 Bräuning, A., Debel, A., Collier, E., Höhnle, S., Mölg, T., Schubert, J.C., Thieroff, B., and Wehrmann, S.:
496 BayTreeNet: Sprechende Bäume als Schnittstelle von Klimadynamik, Dendroökologie und Bildung für
497 nachhaltige Entwicklung in Bayern. *Mitt. der Fränk. Geogr. Ges.*, 67, 177–188, 2022.

498 Braun, V. and Clarke, V.: Using thematic analysis in psychology. *Qual. Res. Psychol.* 3 (2), 77–101,
499 doi:10.1191/1478088706qp063oa, 2006.

500 Boyes, E., and Stanisstreet, M.: The 'Greenhouse Effect': children's perceptions of causes, consequences and
501 cures. *Int. J. Sci. Educ.* 15 (5), 531–552, doi:10.1080/0950069930150507, 1993.

502 Choi, S., Niyogi, D., Shepardson, D. P., and Charusombat, U.: Do Earth and Environmental Science Textbooks
503 Promote Middle and High School Students' Conceptual Development About Climate Change? *Bull. Amer.*
504 *Meteorol. Soc.*, 91 (7), 889–898, doi:10.1175/2009BAMS2625.1, 2010.

505 Clark, R. T. and Brown, S. J.: Influences of Circulation and Climate Change on European Summer Heat Extremes,
506 J. Climate, 26, 9621–9632, doi:10.1175/JCLI-D-12-00740.1, 2013.

507 Collier, E. and Mölg, T.: BAYWRF: a high-resolution present-day climatological atmospheric dataset for Bavaria,
508 Earth Syst. Sci. Data, 12, 3097–3112, <https://doi.org/10.5194/essd-12-3097-2020>, 2020.

509 Davi, N., Pringle, P., Fiondella, F., Lockwood, J., and Oelkers, R.: Online labs to introduce undergraduate students
510 to scientific concepts and practices in tree-ring research, J. Geosci. Educ., 70:1, 73–84,
511 doi:10.1080/10899995.2021.1927567, 2022.

512 Davis, C.C., Lyra, G.M., Park, D.S., Asprino, R., Maruyama, R., Torquato, D., Cook, B.I., and Ellison, A.M.: New
513 directions in tropical phenology. Trends Ecol. Evol., 37(8), 683–693, doi:10.1016/j.tree.2022.05.001, 2022.

514 Debel, A., Meier, W. J. H., and Bräuning, A.: Climate signals for growth variations of *F. sylvatica*, *P. abies*, and
515 *P. sylvestris* in southeast Germany over the past 50 years. Forests, 12, 1433,
516 doi:<https://doi.org/10.3390/f12111433>, 2021.

517 Deslauriers, A., Rossi, S., and Anfodillo, T.: Dendrometer and intra-annual tree growth: What kind of information
518 can be inferred? Dendrochronologia, 25 (2), 113–124, doi:10.1016/j.dendro.2007.05.003, 2007.

519 Dove, J.: Student Teacher Understanding of the Greenhouse Effect, Ozone Layer Depletion and Acid Rain.
520 Environ. Educ. Res., 2 (1), 89–100, doi:10.1080/1350462960020108, 1996.

521 Drew, D., and Downes, G. M.: The use of precision dendrometers in research on daily stem size and wood property
522 variation: A review. Dendrochronologia 27 (2), 159–172, doi:10.1016/j.dendro.2009.06.008, 2009.

523 Dulamsuren, C., Hauck, M., Kopp, G., Ruff, M., and Leuschner, C.: European beech responds to climate change
524 with growth decline at lower, and growth increase at higher elevations in the center of its distribution range
525 (SW Germany). Trees, 31 (2), 673–686, doi:10.1007/s00468-016-1499-x, 2017.

526 Friedrichs, D. A., Trouet, V., Büntgen, U., Frank, D. C., Esper, J., Neuwirth, B., and Löffler, J.: Species-specific
527 climate sensitivity of tree growth in Central-West Germany. Trees, 23 (4), 729–739, doi:10.1007/s00468-009-
528 0315-2, 2009.

529 Herrera-Lormendez, P., Mastrantonas, N., Douville, H., Hoy, A., and Matschullat, J.: Synoptic circulation changes
530 over Central Europe from 1900 to 2100: Reanalyses and Coupled Model Intercomparison Project phase 6, Int.
531 J. Climatol., 42, 4062–4077, doi:10.1002/joc.7481, 2022.

532 James, P. M.: An objective classification method for Hess and Brezowsky Grosswetterlagen over Europe, Theor.
533 Appl. Climatol., 88, 17–42, doi: 10.1007/s00704-006-0239-3, 2007.

534 Kraus, C., Zang, C., and Menzel, A.: Elevational response in leaf and xylem phenology reveals different
535 prolongation of growing period of common beech and Norway spruce under warming conditions in the
536 Bavarian Alps. Eur. J. Forest Res., 135 (6), 1011–1023, doi:10.1007/s10342-016-0990-7, 2016.

537 Lazo J. K., Morss R. E., and Demuth J. L.: 300 Billion served. Sources, Perceptions, Uses, and Values of Weather
538 Forecasts. Bull. Amer. Meteorol. Soc., 90, 785–798, doi:10.1175/2008BAMS2604.1, 2009.

539 Leiserowitz, A., Roser-Renouf, C., Marlon, J., and Maibach, E.: Global Warming’s Six Americas: a review and
540 recommendations for climate change communication. Curr. Opin. Behav. Sci., 42, 97–103,
541 doi:10.1016/j.cobeha.2021.04.007, 2021.

542 Podschuweit, S., and Bernholt, S.: Investigating network coherence to assess students’ conceptual understanding
543 of energy. Educ. Sci., 10(4), 1–20, doi:10.3390/educsci10040103, 2020.

544 Loikith, P. C., Lintner, B. R., and Sweeney, A.: Characterizing Large-Scale Meteorological Patterns and
545 Associated Temperature and Precipitation Extremes over the Northwestern United States Using Self-
546 Organizing Maps. *J. Climate*, 30, 2829–2847, doi:10.1175/JCLI-D-16-0670.1, 2017.

547 Post, P., Truija V., and Tuulik, J.: Circulation weather types and their influence on temperature and precipitation
548 in Estonia, *Boreal Env. Res.*, 7, 281–289, 2002.

549 Psistaki, K., Paschalidou, A. K., and McGregor, G.: Weather patterns and all-cause mortality in England, UK, *Int.*
550 *J. Biometeorol.*, 64, 123–136, doi:10.1007/s00484-019-01803-0, 2020.

551 Riediger, U., and Gratzki, A.: Future weather types and their influence on mean and extreme climate indices for
552 precipitation and temperature in Central Europe, *Meteorol. Z.*, 23 (3), 231–252, doi:10.1127/0941-
553 2948/2014/0519, 2014.

554 Steppe, K., De Pauw, D. J. W., Lemeur, R., and Vanrolleghem, P. A.: A mathematical model linking tree sap flow
555 dynamics to daily stem diameter fluctuations and radial stem growth. *Tree Physiol.*, 26, 257–273, 2006.

556 Steppe, K., De Pauw, D. J. W., Doody, T. M., and Teskey, R. O.: A comparison of sap flux density using thermal
557 dissipation, heat pulse velocity and heat field deformation methods. *Agric. For. Meteorol.*, 150, 1046–1056,
558 doi:10.1016/j.agrformet.2010.04.004, 2010.

559 Steppe, K., Sterck, F., and Deslauriers, A.: Diel growth dynamics in tree stems: linking anatomy and
560 ecophysiology. *Trends Plant Sci.*, 20, 335–343, doi:10.1016/j.tplants.2015.03.015, 2015.

561 Steppe, K., von der Crone, Jonas S., and de Pauw, D. J. W.: TreeWatch.net: A Water and Carbon Monitoring and
562 Modeling Network to Assess Instant Tree Hydraulics and Carbon Status. *Frontiers Plant Sci.*, 7, 993,
563 doi:10.3389/fpls.2016.00993, 2016.

564 Smith, D. M., and Allen, S. J.: Measurement of sap flow in plant stems. *J. Exp. Bot.*, 47 (12), 1833–1844.
565 doi:10.1093/jxb/47.12.1833, 1996.

566 Vandegehuchte, M. W., and Steppe, K.: Sapflow+: a four-needle heat-pulse sap flow sensor enabling nonempirical
567 sap flux density and water content measurements. *The New Phytologist*, 196 (1), 306–317, doi:10.1111/j.1469-
568 8137.2012.04237.x, 2012.

569 Werner, P. C., and Gerstengarbe, F. W.: PIK Report. Katalog der Großwetterlagen Europas (1881-2009), 2010.

570 Witherby, A. E., and Carpenter, S. K.: The rich-get-richer effect: Prior knowledge predicts new learning of domain-
571 relevant information. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, 48(4), 483–498,
572 doi:10.1037/xlm0000996, 2022.

573 Zong, L., Yang, Y., Xia, H., Gao, M., Sun, Z., Zheng, Z., Li, X., Ning, G., Li, Y., and Lolli, S.: Joint occurrence
574 of heatwaves and ozone pollution and increased health risks in Beijing, China: role of synoptic weather pattern
575 and urbanization, *Atmos. Chem. Phys.*, 22, 6523–6538, doi:10.5194/acp-22-6523-2022, 2022.